

# FLUKICIDE® 12.5%

## Broad Spectrum Anthelmintic

To treat worm infestation

### COMPOSITION (each liter contains):

Albendazole ..... 125 g

### DESCRIPTION:

FLUKICIDE® 12.5 % is broad spectrum anthelmintic and effective against egg or larvae of nematodes, cestodes, and trematodes.

### INDICATION:

FLUKICIDE® 12.5 % is recommended to treat worm infestation in cattle, sheep, goat and horse caused by:

- Nematodes
  - Gastrointestinal worms: *Haemonchus* spp., *Chabertia* spp., *Trichostrongylus* spp., *Ostertagia* spp., *Bunostomum* spp., *Cooperia* spp., *Nematodrius* spp., *Oesophagostomum* spp., *Stongyloides* spp., *Oxyuris* spp. etc;
  - Lung worms: *Dictyocaulus* spp.
- Trematodes (*Fasciola* spp., *Dicrocoelium* spp.)
- Cestodes (*Moniezia* spp.)

### DOSAGE AND ADMINISTRATION:

Kinds of Helminth	Animals	Dosage	Note
Nematode, Cestode	Cattle	6 ml / 100 kg of body weight	Administered orally through drinking water or feed  Can be repeated 3-4 month later
	Horse	4 ml / 100 kg of body weight	
	Sheep, Goat	2 ml / 50 kg of body weight	
	Poultry	1 liter for 8,000-16,000 chickens (body weight @ 1.5 kg)	
Trematode	Cattle	8 ml / 100 kg of body weight	Can be repeated 3-4 month later
	Sheep, Goat	3 ml / 50 kg of body weight	

### PRECAUTION:

- Should not be given to cows during the first 45 days of pregnancy and ewes during the first 30 days of pregnancy
- Animal should not be slaughtered until 14 days for cattle and until 10 days for sheep/ goat after the last treatment

### PACKAGING:

- 100 ml (bottle)
- 1 l (jar)

### REG. No.:

DEPTAN RI No. D 0206521 PKC